



Does the Use of a Light Shot Put Modify the Throwing Pattern of Elite Athletes?

Daniel Dinu, Françoise Natta, Cécile Huiban, Nicolas Houel

► To cite this version:

Daniel Dinu, Françoise Natta, Cécile Huiban, Nicolas Houel. Does the Use of a Light Shot Put Modify the Throwing Pattern of Elite Athletes?. *Procedia Engineering*, 2014, 72, pp.92-96. 10.1016/j.proeng.2014.06.019 . hal-01869790

HAL Id: hal-01869790

<https://insep.hal.science/hal-01869790>

Submitted on 6 Sep 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/263712689>

Does the Use of a Light Shot Put Modify the Throwing Pattern of Elite Athletes?

Article in *Procedia Engineering* · December 2014

DOI: 10.1016/j.proeng.2014.06.019

CITATION

1

READS

289

4 authors, including:



Daniel Dinu

Institut National du Sport, de l'Expertise et de la Performance

21 PUBLICATIONS 26 CITATIONS

[SEE PROFILE](#)



Nicolas Houel

Ecole Supérieure d'Ostéopathie

36 PUBLICATIONS 141 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



STUDY OF PUNCHES PERFORMANCE IN BOXE USING INETIAL SENSORS [View project](#)



Study of Lunge Biomechanics in modern fencing sabre, foil, and sword [View project](#)

The 2014 conference of the International Sports Engineering Association

Does the use of a light shot put modify the throwing pattern of elite athletes?

Daniel Dinu^a, Françoise Natta^a, Cecile Huiban^b, Nicolas Houel^c *

^aDépartement de la Recherche, Institut National du Sport, de l'Expertise et de la Performance, 11 Av. du Tremblay, Paris, France

^bDépartement Médical, Institut National du Sport, de l'Expertise et de la Performance, 11 Av. du Tremblay, Paris, France

^cEcole Supérieure d'Ostéopathie-Paris, Cité Descartes, 8, Rue Alfred Nobel, Champs sur Marne, France

Abstract

The aim of this study is to analyse the influence of the weight mass (6 vs 7.26 kg) in kinematics parameters on an expert throwing shot put in rotation, in order to estimate the influence of using a lighter weight to minimise the arm thrower's risk of injury or during recovery training. This study was done on a sample selection of three elite throwers (best performance 16.51 ± 0.81 m) from the French national team, who performed three throws with a 6 Kg weight mass and three others with a 7.26 Kg weight mass. Motion analysis of each throw were recorded using an optoelectronic system (Vicon612) composed of six infrared camcorders and a 3-D modelling of the throwing movement was processed according to the Dapena method. The thrower and weight center of mass velocities were computed for each throwing phase. Friedman Anova analysis showed no significant difference on the subject center of mass and weight velocities (respectively $p = 0.86$ and $p = 0.45$) between both weight mass conditions. No significant shot put mass effect, ($p = 0.86$) on the phases duration in both conditions of weight (6 vs 7.2 kg), were observed. These results suggest that a weight mass lower than 27 % could be used for elite athletes training without a change in velocity of the motion throw. This could be used in order to narrow musculo-skeletal overwork and injury during intensive phases of training and/or to quickly return to specific throwing motion during recovery of the shoulder complex. This should allow for the use of a lighter discus during high level training without kinematical alteration of throwing.

© 2014 Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license

(<http://creativecommons.org/licenses/by-nc-nd/3.0/>).

Selection and peer-review under responsibility of the Centre for Sports Engineering Research, Sheffield Hallam University

Key words: Shot put, Kinematics, Modelling, Throwing technique, Prevention.

* Corresponding author. Tel.: +33(0)141744177; fax: +33(0)141744535

E-mail address: daniel.dinu@insep.fr

1. Introduction

Shot put is a complex movement that associates segment's translation and rotation (Ariel, 1973; Zatsiorsky et al., 1981; Lanka, 2000; McCoy and Gregor, 1984; Hubbard, 1989). The goal is to release the shot put with maximum forward velocity at an angle of approximately forty degrees (Bashian et al., 1982; Hubbard et al., 2001; Linthorne, 2001; Bartonietz, 1996). Nowadays, most of the high level competitors use the spin style whose main purposes is to reach a high rotational speed of the body and to transfer the energy into the shot put, Linthorne (2001). As in discus studies, detailed analysis of the shot put throwing in rotation is facilitated by subdividing the movement into five phases:

- Preparation, double support phase starting from the change in shot put direction at the end of its backward swing and ending when the right foot breaks contact (Δt_{1da}).
- Entry, single support phase which finishes with the left foot breaking contact (Δt_{1sa}).
- Airborne, which finishes with the right foot re-contacting (Δt_s).
- Transition, single support phase which ends as the left foot lands (Δt_{2sa}).
- Delivery, which starts as double support phase and which ends at release of the shot put (Δt_{2da}).

Each phase is separated by “key moments” mostly related to foot contact. Even if the release conditions have been habitually recorded, few studies have focused on the thrower's co-ordination (McCoy and Gregor, 1984; Linthorne, 2001). In order to perform the shot put, the thrower must control a complex motion that is executed in the limited space of the shot put area in agreement with the IAAF rules (IAAF 2011). The best throwers are able to improve their rotation velocity during the delivery phase (McCoy and Gregor, 1984 ; Linthorne, 2001). However, this velocity improvement when associated with the shot put mass can increase shoulders injuries during competition. Edouard et al. (2010) showed that 75% of throwers presented one or more injuries of the throwing arm during their careers. The shoulder was the most commonly injured body part (70%) (Edouard et al., 2010). In order to limit the injury risk and/or for faster recovery, coaches (Cappos, 2014; Wyatt, 2003) recommended training using a lighter shot put (6 vs 7.26 kg). Dinu et al. (2008) have shown similar results in the discus throws. Presently, few studies report the benefits of performing while using a lighter shot put (Judge et al., 2013).

2. Methods

The experiment took place in an indoor hall at the Institut National du Sport, de l'Expertise et de la Performance (Paris, France). Three male subject volunteers (21.33 ± 1.53 years, 128.57 ± 21.83 kg, 190 ± 6 cm, best performance : 16.51 ± 0.81 m) participated in this study. All subjects were high level shot putters from the French national team. All subjects were asked to perform a series of 20 throws using both 6 vs 7.26 kg shot puts, in random order. The throwing distance was measured after each shot put. For each subject, only the three best shot puts (longest distance) in both weight categories were studied. In total 18 throws were studied. Kinematic data were collected by Vicon612 systems (Vicon Motion systems, Oxford, UK). The optoelectronic system (Vicon) consisted of six infrared camcorders. A static calibration around the throwing area was done. A dynamic calibration was performed using a 390 mm stick with two markers moving in the throwing area. The sampling frequency of the Vicon system was 200 Hz. For this experiment, 17 reflective markers were attached on the joints of the athlete's bodies. Reflexive markers were fixed on the head vertex, right and left acromions, lateral humeral epicondyles, ulnar styloids, second metacarpal heads, superior iliac spines, fibula heads, lateral malleolus and second toe tips. The subject's center of mass was computed by the markers' trajectories made by the subject and anthropometric data according to De Leva (1996). In order to determine velocity of the reflective markers, numerical derivation of the kinematics data was performed using the moving average rule. Thrower and weight centre of mass velocities were computed for each throwing phase. Data was collected starting from the preparation stage to the delivery stage. A Friedman Anova was used to compare the significant difference between the velocities and the time of each phase measured between both mass shot puts (6 vs 7.2 kg).

3. Results

The Friedman Anova analysis showed no significant shot put mass effect ($p = 0.45$) on velocities during the different phases of throw in both conditions of weight (6 vs 7.26 kg). In addition, there was no significant shot put mass effect ($p = 0.86$) on the subject center of mass velocities (Table 2) during the different phases of throw in both conditions of weight (6 vs 7.26 kg). Likewise, there was no significant shot put mass effect ($p = 0.86$) on the phases duration in the both conditions of weight (6 vs 7.2 kg).

The kinetics of shot put velocity (6 vs 7.26 kg) during the different phases of the throw is presented on Figure 1 and Table 1, below. After an increasing shot put velocity during the preparation phase, noticeably there is a relative plateau in the shot put velocity during the entry and airborne phases. Velocity began to increase again at the end of transition phase and during delivery. The shot put velocity is maximal at release. The delivery phase has been reported as responsible for 62 % of the release speed of the shot put.




Fig. 1. Comparison of shot put velocity for the two conditions of mass 6 vs 7.2 kg.

Table 1. Kinematics' parameters of shot put.

Kinematics parameters	Time instant & phases	Mean ($\pm$ SD)			
		Shot put 6 kg		Shot put 7.26 kg	
Vcm (m.s⁻¹)	t1da	0,81	$\pm$ 0,13	0,81	$\pm$ 0,15
	t1sa	2,16	$\pm$ 0,17	2,19	$\pm$ 0,16
	ts	2,11	$\pm$ 0,21	2,14	$\pm$ 0,23
	t2sa	1,80	$\pm$ 0,21	1,83	$\pm$ 0,20
	t2da	1,47	$\pm$ 0,39	1,43	$\pm$ 0,36

Table 2. Kinematics' parameters of thrower's center of mass.

Kinematics parameters	Time instant & phases	Mean ($\pm$ SD)					
		Shot put 6 kg			Shot put 7.26 kg		
Vsp (m.s⁻¹)	t1da	2,67	$\pm$	0,20	2,85	$\pm$	0,22
	t1sa	1,84	$\pm$	0,13	1,89	$\pm$	0,27
	ts	1,43	$\pm$	0,11	1,53	$\pm$	0,28
	t2sa	2,79	$\pm$	0,22	2,59	$\pm$	0,95
	t2da	9,83	$\pm$	0,35	9,30	$\pm$	0,21

4. Discussion

The result of this study shows that the subject center of mass and shot put velocities are not disrupted by the weight shot put mass difference (6 vs 7.26 kg) during each phase of the throw. The limitation of this study is related to the relatively small number of subjects studied. However, these athletes are a good representative sample of the best European elite population.

The results demonstrate that the shot put velocities were not influenced by the weight shot put mass difference (6 vs 7.26 kg) during each phase of the throw. The mean shot put velocities are in agreement with the Dapena (1993) results. Furthermore, the results of this study are consistent with previous studies observed on discus thrower (Schluter and Nixdorf, 1984; Schöllhorn, 1989; Dinu et al. 2003, 2008) with different discus weight (1.7 vs 2 kg). Therefore, in order to improve the recovery of the technical level of athletes after injury, it would be beneficial to use a light weight during the training. This could reduce the risk of a new injury related to mechanical load and musculoskeletal stress. According to Ohberg and Alfredson (2004), sub maximal exercise seems to be associated with a more normal tendon structure and no remaining neovascularisation. Training with a light weight shot put would be beneficial for the tendon of the shoulder complex. This could also help prevent shoulder tendon and ligament injuries.

According to the discus throw analysis, it has been observed that the shot put velocity tends to increase at release when athletes used light weight shot puts. This increasing velocity occurred especially during the delivery phase when the athlete threw the light weight shot put (6 kg). This increasing velocity observed during the delivery phase could improve the shot put performance. This tendency is in agreement with previous coaches' recommendations (Cappos, 2014; Wyatt, 2003). However, this tendency has no statistical significant effect on the shot put velocities. Using a similar rotation techniques, Lindsay (1991) and Knicker (1990) observed comparable increasing velocities during the delivery phase in discus throwing analysis. In contrast with the shot put velocities, the subject's center of mass velocities seem much closer during each phase of the throwing motion for both weight shot puts mass. This result is in agreement with previous studies observed on center of mass discus thrower (Dinu et al., 2008).

5. Conclusion

This study showed that the use of a lighter shot put did not influence the shot put and center of mass velocities and phases' durations on elite throwing motion. These results suggest that a weight mass lower than 27 % could be used for elite athlete training without velocities change of the throw's motion. Moreover, this could be used in order to narrow musculo-skeletal overwork and injury during intensive phases of training. It could also be used in order to improve the return to specific throwing motion during recovery of the shoulder complex. Consequently, a lighter discus maybe used for high level training without affecting the kinematical alteration of throwing.

References

- Ariel, G.B., 1973. Computerized biomechanical analysis of the world's best shot-putters. *Track and Field Quarterly Review* 73, 199-206.
- Bartonietz, K., 1996. The energy relationship in rotational and glide shot put techniques. *Modern Athlete and Coach* 34, 7-10.
- Bashian, A., Gavoor, N., and Clark, B., 1982. Some observations on the release in the shotput. *Track and Field Quarterly Review* 82, 12.

- Cappos S. 2014. 10 Glide Shot Put Coaching Tips. <http://digitaltrackandfield.com/glide-shot-put-coaching-tips/>.
- Dinu, D., Natta, F., et Vandewalle, H., 2003. Etude des paramètres cinématiques du lancer de disque : influence du niveau d'expertise dans les différentes phases du lancer. *Science et Motricité* 19, 189-192.
- Dinu, D., Natta, F., Portero, P., Vandewalle, H., 2008. Does the use of a light discus modify the throwing pattern? A study of kinematical and electromyographical data of the throwing arm. *The engineering of sport*, Ed. Springer, pp. 689-694.
- Dapena, J., 1993. An analysis of angular momentum in the discus throw. *ISB 93 Paris I*, 306-307.
- De Leva P., 1996. Adjustment to Zatsiorsky-Seluyanov's segment inertia parameters. *Journal of Biomechanics* 29, 1223-1230.
- International Association of Athletics Federations (IAAF). 2011. Competition rules 2012-2013, in force as from 1st November 2011. Centenary Edition. <http://www.iaaf.org>.
- Edouard P., Depiesse F., Serra J.M. 2010. Throwing arm injuries in high-level athletics throwers. *Science & Sports* 25, 318-322.
- Hubbard, M., 1989. The throwing events in track and field. In: Vaughan CL (Ed.) *Biomechanics of Sport*, Boca Raton, Florida : CRC Press; pp. 213-238.
- Hubbard, M., De Mestre, N.J., Scott J., 2001. Dependence of release variables in the shot put. *Journal of Biomechanics* 34, 449-456.
- Judge L.W., Bellar D., Graig B., Gilreath E., Cappos S., Trasher A. (2013). The influence of post activation potentiation on shot put performance of collegiate throwers. *Journal of Strength Conditionnal Research Epub ahead of print*.
- Knicker, AJ., 1990. Determining factors in discus throwing of top level athletes. In *Proceedings of the First International conference on Techniques in Athletics*. G.-P. Brüggemann and J.K. Rühl, (Ed.); Deutsche Sporthochschule 2, pp. 670-677.
- Lanka, J., 2000. Shot putting. In Zatsiorsky VM (Ed.), *Biomechanics in Sport, Performance Enhancement and Injury Prevention* ; Oxford: Blackwell Science, pp. 435-457.
- Lindsay, M., 1991. Biomechanical analysis of the discus. In *Report on the 1990 Championship in Athletics*, vol. 1.; *The throws* (Ed), Bartlett, R.M; Alsager: Crewe and Alsager College of Higher Education, pp. 47-53.
- Linthorne, N.P., 2001. Optimum release angle in the shot put. *Journal of Sports Sciences* 19, 359-72.
- McCoy, R. & Gregor, P., 1984. Kinematic analysis of elite shot-putters. *Track Technique* 90, 2868-2871.
- Ohberg, L. & Alfredson, H., 2004. Effects on neovascularisation behind the good results with eccentric training in chronic mid-portion Achilles tendinosis? *Knee surgery sports. Traumato Arthrosc* 465-470.
- Schlüter, W. and Nixdorf, E., 1984. Kinematische Beschreibung und Analyse der Diskuswurftechnik. *Leistungssport* 6, 17-22.
- Schöllhorn, W., 1989. Diskuswurf. In *Biomechanik der Sportarten* (edited by K. Willimczik), Reinbek bei Hamburg: Rowohlt Taschenbuch Verlag, pp. 212-19.
- Wyatt P. 2003. Train light throw big. <http://www.canthrow.com/articles/ltimplements.shtml>.
- Zatsiorsky, V.M., Lanka, J.J., Shalmanov, A.A., 1981. Biomechanical analysis of shot putting technique. *Exercise Sport Science Review* 9, 353-89.