



HAL
open science

High-intensity exercise in hypoxia studies shows that training intensity promotes increase in mitochondrial content

Franck Brocherie, Olivier Girard, Grégoire P Millet

► **To cite this version:**

Franck Brocherie, Olivier Girard, Grégoire P Millet. High-intensity exercise in hypoxia studies shows that training intensity promotes increase in mitochondrial content. *The Journal of Physiology*, 2019. hal-02544715

HAL Id: hal-02544715

<https://insep.hal.science/hal-02544715v1>

Submitted on 16 Apr 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Title: High-intensity exercise in hypoxia studies shows that training intensity promotes increase in mitochondrial content

Authors: F. Brocherie¹, O Girard², GP Millet³

¹ Laboratory Sport, Expertise and Performance (EA 7370), French Institute of Sport (INSEP), Paris, France.

² Murdoch Applied Sports Science Laboratory, College of Science, Health, Engineering and Education (SHEE), Murdoch University, Perth, Australia.

³ Institute of Sport Sciences (ISSUL), University of Lausanne, Switzerland.

Corresponding author:

Dr. Franck Brocherie

Laboratory Sport, Expertise and Performance (EA 7370), French Institute of Sport (INSEP)

11, Avenue du Tremblay, 75012 Paris, France

Ph. +33 (0)1 41 74 43 54

Email franck.brocherie@insep.fr

Comment-only count: 201 words

Competing interests: None declared.

Each side of this cross-talk ^{1,2} properly defended their views. Another way to shed light on the importance of training intensity or volume may come from studies on exercise in hypoxia through the interdependence of environmental oxygen availability and mitochondrial oxidative phosphorylation.

Hypoxic exercise can elicit specific transcriptional responses [through the upregulation of hypoxia-inducible factor 1] leading to skeletal muscle adaptations³. Interestingly, high-intensity training in hypoxia induces higher increase in total mitochondrial volume density and capillary length density than low-intensity training in hypoxia ^{4,5}. It has been then postulated that exercise intensity in hypoxia *per se* modulates muscle molecular mechanisms of oxygen homeostasis ⁶. In partial agreement, performing maximal-intensity exercise training in hypoxia (*i.e.*, repeated-sprint training in hypoxia, RSH) *versus* normoxia upregulates genes involved in oxygen signaling, oxygen carrying and pH regulation but not in mitochondrial biogenesis in moderately-trained male cyclists ⁷. In team-sport athletes, however, superimposing RSH to 14-d of hypoxic residence (200 h) elicits higher oxygen signaling and oxygen carrying as well as mitochondrial biogenesis and mitochondrial metabolism compared with similar normoxic intervention ⁸. Altogether, these recent studies on high-intensity hypoxic exercise tend to support that '*training intensity is of primary importance for increasing human skeletal muscle mitochondrial content*'.

Competing interests

None declared.

References

1. Bishop DJ, Botella J, Granata C. CrossTalk opposing view: Exercise training volume is more important than training intensity to promote increases in mitochondrial content. *J Physiol* 2019.
2. MacInnis MJ, Skelly LE, Gibala MJ. CrossTalk proposal: Exercise training intensity is more important than volume to promote increases in human skeletal muscle mitochondrial content. *J Physiol* 2019.
3. Faiss R, Girard O, Millet GP. Advancing hypoxic training in team sports: from intermittent hypoxic training to repeated sprint training in hypoxia. *Br J Sports Med* 2013;47 Suppl 1:i45-i50.
4. Vogt M, Puntschart A, Geiser J, Zuleger C, Billeter R, Hoppeler H. Molecular adaptations in human skeletal muscle to endurance training under simulated hypoxic conditions. *J Appl Physiol* 2001;91:173-82.
5. Geiser J, Vogt M, Billeter R, Zuleger C, Belforti F, Hoppeler H. Training high--living low: changes of aerobic performance and muscle structure with training at simulated altitude. *Int J Sports Med* 2001;22:579-85.
6. Hoppeler H, Vogt M. Muscle tissue adaptations to hypoxia. *J Exp Biol* 2001;204:3133-9.
7. Faiss R, Leger B, Vesin JM, et al. Significant molecular and systemic adaptations after repeated sprint training in hypoxia. *PLoS One* 2013;8:e56522.
8. Brocherie F, Millet GP, D'Hulst G, Van Thienen R, Deldicque L, Girard O. Repeated maximal-intensity hypoxic exercise superimposed to hypoxic residence boosts skeletal muscle transcriptional responses in elite team-sport athletes. *Acta Physiol (Oxf)* 2018;222:e12851.